

No.

200200115



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Svalof Weibull A B

Whereas, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

PEA, FIELD

'SW Circus'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C. this twenty ninth day of April, in the year two thousand three.


Attest:

P. M. Jakob

Commissioner
Plant Variety Protection Office
Agricultural Marketing Service

A. Freeman

Secretary of Agriculture



U.S. DEPARTMENT OF AGRICULTURE
 AGRICULTURAL MARKETING SERVICE
 SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE
 (Instructions and information collection burden statement on reverse)

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

1. NAME OF OWNER SVALOF WEIBULL AB		2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME SW 975504	3. VARIETY NAME SW CIRCUS
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) 5-268 81 SVALOV, SWEDEN		5. TELEPHONE (include area code) 46-418-667000	FOR OFFICIAL USE ONLY 200200115
		6. FAX (include area code) 46-418-667100	
7. IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) CORPORATION	8. IF INCORPORATED, GIVE STATE OF INCORPORATION SWEDEN	9. DATE OF INCORPORATION 1993	FILING DATE MARCH 15, 2002

10. NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THIS APPLICATION. (First person listed will receive all papers) SVALOF WEIBULL LTD. 2-411 DOWNEY RD. SASKATOON, SK CANADA S7N 4L8	FILING AND EXAMINATION FEES: \$ 2705.00 DATE 3/15/02 CERTIFICATION FEE: \$ 320.00 DATE 5/22/02
---	---

11. TELEPHONE (Include area code) 306-477-5230	12. FAX (Include area code) 306-477-5239	13. E-MAIL howard.lowe@swseed.se	14. CROP KIND (Common Name) Pisum sativum (Field pea)
--	--	--	---

18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on reverse) a. <input checked="" type="checkbox"/> Exhibit A. Origin and Breeding History of the Variety b. <input checked="" type="checkbox"/> Exhibit B. Statement of Distinctness c. <input checked="" type="checkbox"/> Exhibit C. Objective Description of Variety d. <input checked="" type="checkbox"/> Exhibit D. Additional Description of the Variety (Optional) e. <input checked="" type="checkbox"/> Exhibit E. Statement of the Basis of the Owner's Ownership f. <input checked="" type="checkbox"/> Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties, verification that tissue culture will be deposited and maintained in an approved public repository) g. <input checked="" type="checkbox"/> Filing and Examination Fee (\$2,705), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office)	19. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? See Section 83(a) of the Plant Variety Protection Act <input checked="" type="checkbox"/> YES (If "yes", answer items 20 and 21 below) <input type="checkbox"/> NO (If "no," go to item 22)
	20. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF CLASSES? <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO IF YES, WHICH CLASSES? <input checked="" type="checkbox"/> FOUNDATION <input checked="" type="checkbox"/> REGISTERED <input checked="" type="checkbox"/> CERTIFIED
	21. DOES THE OWNER SPECIFY THAT THE CLASSES BE LIMITED AS TO NUMBER OF GENERATIONS? <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO IF YES, SPECIFY THE NUMBER 1, 2, 3, etc. 3 FOUNDATION 2 REGISTERED 1 CERTIFIED (If additional explanation is necessary, please use the space indicated on the reverse.)

22. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U. S. OR OTHER COUNTRIES? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.)	23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO IF YES, GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space indicated on reverse.) MAH 5-6-2002
---	--

24. The owners declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate.

The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act.

Owner(s) is(are) informed that false representation herein can jeopardize protection and result in penalties.

SIGNATURE OF OWNER Howard K. Lowe		SIGNATURE OF OWNER	
NAME (Please print or type) DR. HOWARD K. LOWE		NAME (Please print or type)	
CAPACITY OR TITLE CDN. RESEARCH DIRECTOR, V.P.	DATE MARCH 13/02	CAPACITY OR TITLE	DATE

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,705 \$320 filing fee and \$2,385 examination fee, payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. **DO NOT** use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$320 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office

Telephone: (301) 504-5518

FAX: (301) 504-5291

Homepage: <http://www.ams.usda.gov/science/pvp.htm>

ITEM

- 8a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method; (2) the details of subsequent stages of selection and multiplication; (3) evidence of uniformity and stability; and (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 8b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
- (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 8c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 8d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 8e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
9. If "Yes" is specified (*seed of this variety be sold by variety name only, as a class of certified seed*), the applicant **MAY NOT** reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See *Regulations and Rules of Practice, Section 97.103*).
1. See Section 83 of the Act for the Contents and Term of Plant Variety Protection.
 2. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
 3. See Section 5.5 of the Act for instructions on claiming the benefit of an earlier filing date.

1. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

2. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

3. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 690-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

STD-470 (2-99) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces STD-470 (6-98) which is obsolete.

March 11, 2002

CONFIDENTIAL

VARIETY "SW CIRCUS" expt. designation: SW 975504

Origin and Breeding:

SW 975504 was developed by Svalöf Weibull AB, Svalöv, Sweden. The variety is derived from the cross 9211-9-5 x Carneval. The final cross was made in 1994.

9211-9-5 pedigree = Carneval x Sv 92536

Sv 92536 pedigree = LW 8411-1 X(RIGEL X SV U 21421)

Please be advised that LW 8411 should read 'SW 8411' (this was a typo on the application) and it's parentage is Solara x Capella. 'SV U 21421' was marketed in Sweden under the name Fjord.

SW 975504 was developed by Svalof Weibull AB, Svalov, Sweden. This semi-leafless variety originates from the cross (see confidential pedigree information package). The original cross was done in 1994. The breeding method was a pedigreed method and the variety originates from a single plant selection in the F₄. Selection was made for yield, semi-leafless trait, earliness, straw stiffness, and good yellow colour. Breeder seed was bulked in F₅ generation.

Statement of Uniformity and Stability

SW 975504 is uniform and stable. No offtypes. Stability has been observed in more than two seasons.

Methods of maintaining the variety.

SW 975504 is maintained from breeder seed. Breeder seed will be maintained by Svalöf Weibull AB, Sweden and Svalof Weibull Ltd. Saskatoon, SK, Canada.

No variants were observed.

Variety: SW CIRCUS (SW 975504) Field Pea

Exhibit B: Statement of Distinctness

SW 975504 is a distinct variety, possible to distinguish from Carneval and Majoret which are the most similar varieties known to us. SW 975504 differs from Majoret by having yellow seed and a blunt pod. SW 975504 differs from Carneval by having 2 days earlier maturity. SW 975504 also has larger stipules and higher thousand grain weight than Carneval.

SW CIRCUS differs from Majoret by having blunt pods and yellow seed whereas Majoret has pointed pods and green seed.

We have shown 4 significant differences between SW CIRCUS and Carneval. They are stem vine length, stipule length, stipule width and pod width. Please find attached a trait comparison chart (based on the T-test), trial location information and the raw data for SW CIRCUS and Carneval from 2000 and 2001 in Svalov, Sweden as requested.

Trait comparison between two pea varieties, SW CIRCUS and Carneval (T-test)
****comparisons based on data taken in the years 2000 and 2001 in Svalov, Sweden**

Trait	T-value	P-range
Analysis on the data from Svalov 2000		
Stem vine length	6.47	< 0.001
Stipule length	6.84	< 0.001
Stipule width	7.08	< 0.001
Pod width	3.43	0.001 < P < 0.002

Analysis on the data from Svalov 2001		
Stem vine length	2.86	0.005 < P < 0.01
Stipule length	3.93	< 0.001
Stipule width	3.21	0.002 < P < 0.005
Pod width	4.01	P < 0.001

	2000		2001	
	<u>SW Circus</u>	<u>SW Carneval</u>	<u>SW Circus</u>	<u>SW Carneval</u>
Stem Vine Length (cm):	53.6	45.04	60.08	56.56
Stipule Length (mm):	53.88	48	65.56	58.4
Stipule Width (mm):	25.4	21.72	29.2	26.12
Pod Width (mm):	13.76	12.8	12.96	13.88

PVP Application for SW CIRCUS Appl. #200200115

Year	2000 SW CIRCUS	2000 Carneval	2001 SW CIRCUS	2001 Carneval
Trial Location	Svalov, Sweden	Svalov, Sweden	Svalov, Sweden	Svalov, Sweden
Planting Date	28-Apr-00	28-Apr-00	12-Apr-01	12-Apr-01
Harvest Date	17-Aug-00	17-Aug-00	15-Aug-01	15-Aug-01
Date of Vine Length Note	20-Jun-00	20-Jun-00	11-Jul-01	11-Jul-01
Date of Stipule Length Note	25-Jun-00	25-Jun-00	22-Jun-01	22-Jun-01
Date of Stipule Width Note	25-Jun-00	25-Jun-00	22-Jun-01	22-Jun-01
Date of # Nodes Note	15-Jul-00	15-Jul-00	11-Jul-01	11-Jul-01
Date of Pod Length Note	14-Aug-00	14-Aug-00	15-Aug-01	15-Aug-01
Date of Pod Width Note	14-Aug-00	14-Aug-00	15-Aug-01	15-Aug-01
Date of # Ovules Note	14-Aug-00	14-Aug-00	15-Aug-01	15-Aug-01
Plant Spacing	5cm between seeds rows 10cm apart	5cm between seeds rows 10cm apart	5cm between seeds rows 10cm apart	5cm between seeds rows 10cm apart
Sample Size	25	25	25	25

200200115

PVP Application for SW CIRCUS Appl. #200200115
2000 Svalov, Sweden-SW CIRCUS

	Stem Vine Length	Stipule Length	Stipule Width	Pod Width
	48	55	28	14
	50	54	24	14
	54	51	22	14
	52	53	26	13
	55	56	26	13
	51	53	25	14
	50	54	26	14
	55	55	25	13
	48	50	25	15
	57	50	26	13
	59	57	27	14
	58	58	27	14
	56	56	26	14
	52	58	28	14
	60	58	28	14
	57	53	25	14
	50	54	26	14
	55	56	27	13
	52	58	28	13
	54	56	24	15
	55	53	24	13
	53	50	24	13
	54	48	23	14
	53	51	22	14
	52	50	23	14
AVG	53.6	53.88	25.4	13.76
ST DEV	3.201562119	2.976575213	1.825741858	0.5972158
n=25				

200200115

PVP Application for SW CIRCUS Appl. #200200115

2000 Svalov, Sweden-Carneval

	Stem Vine Length	Stipule Length	Stipule Width	Pod Width
	39	45	20	14
	48	47	20	12
	35	47	21	14
	45	50	22	12
	40	48	23	13
	49	46	21	13
	36	45	22	14
	50	52	20	13
	56	50	22	13
	49	50	21	13
	41	55	25	12
	47	45	19	13
	45	48	23	13
	40	50	22	14
	40	48	22	13
	52	46	21	14
	50	48	21	12
	45	50	23	11
	43	49	20	10
	47	48	22	13
	48	46	24	11
	46	48	22	15
	43	45	21	13
	48	46	23	13
	44	48	23	12
AVG	45.04	48	21.72	12.8
ST DEV	5.020624131	2.415229458	1.4	1.118034
n=25				

200200115

PVP Application for SW CIRCUS Appl. #200200115

2001 Svalov, Sweden-SW CIRCUS

	Stem Vine Length	Stipule Length	Stipule Width	Pod Width
	50	61	29	13
	60	62	30	14
	55	54	24	13
	56	71	32	13
	60	59	27	13
	56	65	25	12
	55	68	30	12
	55	64	26	14
	60	60	24	13
	58	72	31	13
	61	62	30	13
	61	70	30	12
	63	66	25	13
	62	43	34	12
	63	66	28	12
	66	73	37	14
	58	65	30	13
	61	64	28	13
	57	72	30	13
	63	74	28	13
	65	66	30	14
	60	66	28	13
	62	68	28	13
	63	70	31	13
	72	78	35	13
AVG	60.08	65.56	29.2	12.96
ST DEV	4.452714528	7.112664761	3.214550254	0.6110101
n=25				

200200115

PVP Application for SW CIRCUS Appl. #200200115

2001 Svalov, Sweden-Carneval

	Stem Vine Length	Stipule Length	Stipule Width	Pod Width
	59	63	33	13
	53	61	29	12
	59	59	27	13
	59	65	27	13
	55	58	27	14
	57	58	29	14
	57	60	27	15
	57	56	22	13
	53	62	27	14
	53	66	32	14
	61	57	25	14
	56	55	24	14
	50	56	25	13
	54	54	21	14
	60	63	25	15
	49	54	23	15
	55	53	27	14
	59	57	25	15
	58	51	25	14
	59	60	25	13
	60	57	26	14
	60	65	25	15
	56	58	29	13
	55	56	22	14
	60	56	26	15
AVG	56.56	58.4	26.12	13.88
ST DEV	3.241398875	3.958114029	2.847806173	0.8326664
n=25				

UNITED STATES DEPARTMENT OF AGRICULTURE
 AGRICULTURAL MARKETING SERVICE
 Science Division
 NATIONAL AGRICULTURAL LIBRARY
 BELTSVILLE, MARYLAND 20705
 OBJECTIVE DESCRIPTION OF VARIETY
 PEA (*PISUM SATIVUM*)

EXHIBIT C
(Pca)

NAME OF APPLICANT(S) SVALOF WEIBULL AB	VARIETY NAME OR TEMPORARY DESIGNATION SW CIRCUIS (SW 975504)
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) S-268 81 SVALOF, SWEDEN	FOR OFFICIAL USE ONLY
	PVPO NUMBER 200200115

Place the appropriate number that describes the varietal character in the boxes below.

Place a zero in first box (e.g. or) when number is either 99 or less or 9 or less.

1. TYPE:

 1 - GARDEN 2 - FIELD 3 - EDIBLE-PODDED

2. MATURITY:

<input type="text" value="16"/> Node number of first bloom:	<input type="text" value=""/> <input type="text" value=""/> <input type="text" value=""/> No. of days to processing	<input type="text" value=""/> <input type="text" value=""/> <input type="text" value=""/> Heat Units
<input type="text" value=""/> <input type="text" value=""/> No. of days Earlier than . . .	<input type="text" value=""/> 1 - ALASKA WR 2 - THOMAS LAXTON WR 3 - LITTLE MARVEL	
<input type="text" value=""/> <input type="text" value=""/> No. of days Later than . . .	<input type="text" value=""/> 4 - WANDO 5 - ALDERMAN WR 6 - AUSTRIAN WINTER	

3. PLANT HEIGHT:

<input type="text" value=""/> <input type="text" value="7"/> <input type="text" value="0"/> CM. HIGH	
<input type="text" value=""/> <input type="text" value=""/> Cm. Shorter than . . .	<input type="text" value=""/> 1 - ALASKA WR 2 - THOMAS LAXTON WR 3 - LITTLE MARVEL
<input type="text" value=""/> <input type="text" value=""/> Cm. Taller than . . .	<input type="text" value=""/> 4 - WANDO 5 - ALDERMAN WR 6 - AUSTRIAN WINTER

4. VINE:

<input checked="" type="text" value="2"/> Habit: 1 = DETERMINATE 2 = INDETERMINATE	<input checked="" type="text" value="2"/> Stockiness: 1 = SLIM (Alaska) 3 = HEAVY (Alderman) 2 = MEDIUM (Thomas Laxton WR)
<input checked="" type="text" value="2"/> Branching: 1 = NONE (Alaska) 2 - 1-2 BRANCHES (Little Marvel) 3 = MORE THAN 2 BRANCHES (Dwarf Gray Sugar)	
<input type="text" value="1"/> Internodes: 1 = STRAIGHT 2 = ZIG ZAG	<input type="text" value="2"/> <input type="text" value="2"/> NUMBER OF NODES

5. LEAFLETS: not present

<input type="text" value=""/> Color: 1 = LIGHT GREEN (Alaska WR) 2 = MED. GREEN (Thomas Laxton WR) 3 = DARK GREEN (Alderman) 4 = OTHER (Specify)	
<input type="text" value=""/> Wax: 1 = NONE 2 = LIGHT 3 = MEDIUM 4 = HEAVY	<input type="text" value=""/> 1 = NOT MARBLED 2 = MARBLED (Alaska)
<input type="text" value=""/> Number of leaflet pairs: 1 = NOT PAIRED 2 = ONE 3 = TWO 4 = THREE OR MORE	

6. STIPULES:

<input checked="" type="text" value="2"/> 1 = LACKING 2 = PRESENT	<input type="text" value="1"/> 1 = NOT CLASPING 2 = CLASPING
<input checked="" type="text" value="2"/> 1 = NOT MARBLED 2 = MARBLED	<input type="text" value=""/> Size (Compared with leaflets): 1 = SMALLER 2 = SAME 3 = LARGER
<input type="text" value=""/> Color (Compared with leaflets): 1 = LIGHTER 2 = SAME 3 = DARKER	

7. FLOWER COLOR:

<input type="text" value=""/> VENATION	<input type="text" value="1"/> STANDARD	<input type="text" value="1"/> WING	<input type="text" value="1"/> KEEL	1 = WHITE 2 = GREENISH 3 = LAVENDER 4 = PURPLE 5 = RED 6 = OTHER (Specify)
--	---	-------------------------------------	-------------------------------------	---

8. PODS:

☒ 2 Shape: 1 = STRAIGHT 2 = SLIGHTLY CURVED ☐ 2 End: 1 = POINTED (Alderman) 2 = BLUNT (Alaska)
☐ 3 CURVED
☒ 2 Color: 1 = LIGHT GREEN (Alaska WR) 2 = MEDIUM GREEN 3 = DARK GREEN (Alderman)
☐ 4 = OTHER (Specify) _____
☐ 1 Surface: 1 = SMOOTH 2 = ROUGH ☒ 2 Surface: 1 = SHINY 2 = DULL
☒ 3 Borne: 1 = SINGLE 2 = DOUBLE 3 = SINGLE AND DOUBLE 4 = SINGLE, DOUBLE, & TRIPEE
☐ 5 = DOUBLE & TRIPLE 6 = TRIPLE 7 = OTHER (Specify) _____
☐ 7 CM. LENGTH ☐ 1 ☐ 3 MM. WIDTH (Between sutures) ☐ 7 NO. SEEDS PER POD

9. SEEDS (95--100 Tenderometer):

n/a

☐ Color: 1 = LIGHT GREEN 2 = GREEN 3 = DARK GREEN 4 = OTHER (Specify) _____
 Seive: % ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ AVERAGE

SEEDS (Dry, Mature):

☒ 4 Shape: 1 = FLATTENED 2 = ANGULAR 3 = OVAL 4 = ROUNDED
☐ 1 Surface: 1 = SMOOTH 2 = DIMPLED ☒ 2 Surface: 1 = SHINY 2 = DULL
☐ 3 WRINKLED
☐ 1 Color Pattern: 1 = MONOCOLOR 2 = MOTTLED 3 = STRIPED 4 = DOTTED
☐ 1 Primary Color: 1 = CREAMY-WHITE 2 = CREAM & GREEN 3 = LIGHT GREEN 4 = MEDIUM GREEN
☐ 5 = DARK GREEN 6 = BLUE-GREEN 7 = YELLOW 8 = BROWN 9 = RED
☒ Secondary Color: 10 = GRAY 11 = BLACK
☐ 1 Hilum Floor Color: 1 = WHITE 2 = TAN ☒ 2 Cotyledon Color: 1 = GREEN 2 = YELLOW 3 = ORANGE
☐ 3 = BLACK

☒ 2 ☒ 4 GRAMS PER 100 SEEDS
 MAH 5-6-02

10. DISEASE: (0 = Not Tested; 1 = Susceptible; 2 = Resistant)

☐ 0 FUSARIUM WILT ☐ 0 NEAR-WILT ☐ 0 DOWNY MILDEW
☐ 1 ASCOCHYTA BLIGHT ☐ 1 POWDERY MILDEW ☐ 0 BACTERIAL BLIGHT
☐ 0 MOSAIC ☐ 0 PEA ENATION MOSAIC ☐ 0 YELLOW BEAN MOSAIC
☐ OTHER (Specify) _____

11. INSECT: (0 = Not Tested; 1 = Susceptible; 2 = Resistant)

☐ 0 APHIDS ☐ OTHER (Specify) _____

12. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED

CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Leafiness	Carneval	Fresh Seed Color	
Leaf Color	majoret	Mature Seed Color	Carneval
Pod Color	Carneval	Seed Shape	Karita
Pod Shape	Carneval	Plant Height	majoret

COMMENTS:

200200115

TABLE OF CONTENTS

A.	ABOUT THE OBJECTIVE DESCRIPTION FORM.....	2
B.	CHARACTERISTICS	2
C.	LEGEND	2
D.	PEA OBJECTIVE DESCRIPTION	3
E.	APPENDICES: ILLUSTRATIONS & METHODS	12

A. ABOUT THE OBJECTIVE DESCRIPTION FORM

This objective description form is designed as an aid for the identification of field pea varieties to provide sufficient information for pedigreed seed crop inspection and variety verification purposes. Companion documents include the "Variety Registration Application Form" and the "Procedures for the Registration of Crop Varieties in Canada", both of which are available from the Variety Registration Office.

This objective description form lists characteristics to be used as the basis for developing the description of field pea varieties. It is recommended that the form be completed in as much detail as possible to ensure that an accurate description of the variety be on record. Uniformity and stability must be sufficient to ensure that the genetic purity of the variety has not been compromised during the development of the variety or during the seed multiplication process. However, accurate information on variability within the variety is essential for distinguishing between variants and off-types during the seed multiplication process.

Information on this document may be accessible or protected as required under the provisions of the *Access to Information Act*. Information that could cause you or your organization injury if released is protected from disclosure as defined in Section 20 of the *Access to Information Act*.

B. TEST GUIDELINES

1. The candidate variety **must** be described for all characteristics designated on the form with the pound symbol (#).
2. A rating system of 1-9 provides a scale for describing most characteristics in this form. To rate characteristics, select a value that best corresponds to the state indicated. Characteristics may be rated with intermediate values where the characteristic grades gradually from one extreme to another. For example, where the states for a characteristic are described as: small (3), medium (5), large (7); other values of 1, 2, 4, 6, 8 or 9 may be selected.
3. Each characteristic on this form has been arranged in a tabular format allowing the candidate variety (CV) and up to four reference/check varieties (R1 to R4) to be described. Information on reference varieties is useful but **not** required for variety registration. The reference varieties must be registered for sale in Canada.

C. LEGEND

- (#) Characteristics that must always be included when completing the objective description form for variety registration, except when the state of expression of a preceding characteristic renders this impossible.
- (+) Indicates an illustration or method for this trait is in the appendix.

CV Candidate variety: **SW 975504**

RI - R4 Reference or check varieties

RI Carneval R3 _____

R2 Majoret R4 _____

D. PEA OBJECTIVE DESCRIPTION

Applicant (name and address):

Bonis & Company208 St. David StreetLindsay, ONK9V 5Z4Telephone: (705) 324-0544 Fax: (705) 324-2550**1.0 CLASSIFICATION (#)**1.1 Botanical name: *Pisum sativum* L.

1.2 Type: ☐ 1. Field - green
☒ 2. Field - yellow

1.2 Proposed variety denomination (name): SW CIRCUS**2.0 PLANT CHARACTERISTICS**

2.1 Plant: growth habit

(#)		CV	R1	R2	R3	R4
Determined (bush type)	1	9	9	9		
Indeterminate (tall type)	9					

2.2 Plant: height (observe when 30% of plants have one flower open)

(#)						
Short (< 25 cm)	3	5	5	5		
Medium (25-50 cm)	5					
Tall (> 50 cm)	7					

2.3 Plant: foliage colour (observe at flowering)

Yellow green	1	2	2	2		
Green	2					
Blue or dark green	3					

3.0 STEM CHARACTERISTICS

3.1 Stem: fasciation

(+)						
Absent	1	1	1	1		
Present	9					

3.2 Stem: vine length (observe after flowering when pods are fully swollen)

(*) (+)

CV

R1

R2

R3

R4

Short (50 – 70 cm)	3	5	5	5		
Medium (90 – 115 cm)	5					
Long (130 – 150 cm)	7					

3.3 Stem: number of nodes up to and including first flowering node (observe at harvest, include the two scale nodes)

(+) (+)

Few	3					
Medium	5					
Many	7					

3.4 Shape of internodes

Straight	1					
Curved	9					

4.0 LEAF CHARACTERISTICS**4.1 Leaf: presence of leaflets**

(*) (*)

Leafed	1	2	2	2		
Semi-leafless	2					
Leafless	3					

4.2 Leaf: average maximum number of leaflets (observe any time after stipules at seventh node are fully opened)

(+) (+)

Four	1					
Six	2					
Eight	3					

4.3 Leaf: size (observe at second fertile node)

Small	3					
Medium	5					
Large	7					

4.4 Leaf: shape (observe at second fertile node)

Elliptic	1					
Ovate	9					

4.5 Leaf: waxiness of leaves and stipules

Absent	1	9	9	9		
Present	9					

4.6 Leaf: dentation (observe over whole plant)

(*)

CV

R1

R2

R3

R4

Absent	1					
Present	9					

4.7 Leaf: degree of dentation

(+)

Very weak	1					
Weak	3					
Medium	5					
Strong	7					
Very strong	9					

4.8 Leaf: apex of leaflet

Pointed	3					
Rounded	5					
Truncate	7					
Retuse	9					

5.0 STIPULE CHARACTERISTICS**5.1** Stipule: development (observe any time after stipules at seventh node are fully opened)

(*) (+)

Rudimentary	1	2	2	2		
Normal	2					

5.2 Stipule: size

(+)

Small	3	3	3	3		
Medium	5					
Large	7					

5.3 Stipule: shape

Elliptic	1					
Broadly elliptic	9					

5.4 Stipule: colouration

Absent	1					
Present	9					

5.5 Stipule: marbling (before flowering, on 2 nodes below the first fertile node)

(#)(+)

Absent	1	9	9	9		
Present	9					

200200115

5.6 Stipule: maximum density of marbling**(#)(+)****CV****R1****R2****R3****R4**

Very sparse	1	2	5	3		
Sparse	3					
Medium	5					
Dense	7					
Very dense	9					

6.0 FLOWERING CHARACTERISTICS**6.1 Time of flowering (observe when approximately 30% of plants have one flower open, avoid recording early flowering variants)****(#)**

Early	3	5	5	5		
Medium	5					
Late	7					

6.2 Maximum number of flowers per node (non-fasciated varieties only, observe when highest nodes show signs of producing flowers which do not develop beyond the bud stage)

One	1	2	2	2		
One to two	2					
Two	3					
Two to three	4					
Three	5					
Three to four	6					
More than four	7					

6.3 Flower: colour of wing**(#)**

White	1	1	1	1		
Greenish	2					
Pink	3					
Purple	4					
Dark red	5					
Other: _____	6					

6.4 Flower: shape of wing**(#)**

Round	1	1	1	1		
Notched	9					

6.5 Flower: colour of standard**(#)**

White	1	1	1	1		
Greenish	2					
Pink	3					
Reddish purple	4					
Other: _____	5					

6.6 Flower: size of standard

		CV	R1	R2	R3	R4
Small	3					
Medium	5					
Large	7					

6.7 Flower: shape of base of standard

(#)(+)

Raised (V-shaped)	3	7	9	7		
Level (straight)	5					
Arched (2 lobes)	7					
Strongly arched	9					

6.8 Flower: apex of calyx lobe (observe at second flowering node)

Acuminate	1					
Pointed	2					
Rounded	3					

7.0 POD CHARACTERISTICS**7.1 Pod: length (observe at first flowering node)**

(#)

Short	3	71.0 mm	69.3 mm	76.5 mm		
Medium	5					
Long	7					

7.2 Pod: width (observe at first flowering node)

(#)

Narrow	3	13.0 mm	13.1 mm	13.8 mm		
Medium	5					
Broad	7					

7.3 Pod: parchment (observe when pods are dry and papery)

(#)(+)

Absent or partially present	1	9	9	9		
Entirely present	9					

7.4 Pod: curvature (observe when pods fully swollen)

(#)

Absent	1	1	3	5		
Weak	3					
Medium	5					
Strong	7					
Very strong	9					

7.5 Pod: type of curvature (observe when pods are fully swollen)

(+)

Towards ventral part	1					
Straight	2					
Towards dorsal part	3					

7.6 Pod: shape of distal part (observe when pods fully swollen)**(+)****CV****R1****R2****R3****R4**

Pointed	1	9	9	1		
Blunt	9					

7.7 Pod: colour (observe when pods fully swollen)**(#) (+)**

Yellow	1	2	2	2		
Green	2					
Blue green	3					
Purple	4					
Other: _____	5					

7.8 Pod: number of ovules/seeds (observe at second fertile node when ovules/seeds are partially developed)

Few	3					
Medium	5					
Many	7					

7.9 Pod: colour of immature seed (observe when seed is firm, before starchy to taste)**(#)**

Light green	1	1	1	1		
Dark blue-green	9					

8.0 SEED CHARACTERISTICS: (+) (observe dry seed)**8.1 Seed: shape of starch grain****(+)**

Simple	1	1	1	1		
Compound	9					

8.2 Seed: colour of cotyledon**(#) (+)**

Green	1	2	2	1		
Yellow	2					
Red	3					

8.3 Seed: black colour of hilum**(#) (+)**

Absent	1	1	1	1		
Present	9					

8.4 Seed: shape**(#) (+)**

Spherical	1	1	1	1		
Irregular	9					

8.5 Seed: wrinkling of cotyledon
(+)

Absent	1	1	1	1		
Present	9					

8.6 Seed: size
(#) (+)

		CV	R1	R2	R3	R4
Small	3	3	3	5		
Medium	5					
Large	7					

8.7 Seed: weight (grams per 1000 seed)
(#)

Weight in grams	240	235	260		
-----------------	-----	-----	-----	--	--

8.8 Time of maturity (observe hard, dry seed)
(#)

Early	3	3	5	5		
Medium	5					
Late	7					

9.0 QUALITY CHARACTERISTICS

9.1 Protein content
(#)

Percentage	21.1	20.8	21.5		
------------	------	------	------	--	--

9.2 Cooking quality (describe) eg. Colour, granulation, viscosity

10.0 REACTION TO DISEASES

- 0 - not tested
 1 - resistant
 3 - moderately resistant
 5 - moderately susceptible
 7 - susceptible
 9 - highly susceptible

		CV	R1	R2	R3	R4
10.1	Seedling blight, root rot and wilt <i>Aphanomyces euteiches</i> <i>Fusarium oxysporum</i> f.sp. <i>pisi</i> <i>Fusarium</i> spp. <i>Pythium</i> spp.					
10.2 (#)	Mycosphaerella blight and ascochyta foot rot <i>Mycosphaerella pinodes</i> <i>Phoma medicaginis</i> var. <i>Pinodella</i>	7	7	5		
10.3	Ascochyta leaf and pod spot <i>Ascochyta pisi</i>					
10.4	Downy mildew <i>Peronospora viciae</i>					
10.5 (#)	Powdery mildew <i>Erysiphe polygoni</i>	5	3	7		
10.6	Bacterial blight <i>Pseudomonas syringae</i> pv. <i>pisi</i>					
10.7	Bean yellow mosaic virus					
10.8	Septoria leaf blotch <i>Septoria pisi</i>					
10.9	Other (specify) _____ _____ _____					

11.0 Describe chemical characteristics that aid in the identification of the candidate variety, eg. electrophoresis. Please attach data and the corresponding protocol.

- 12.0** Describe any deviant plants, including both variants and off-types observed during seed increase of the candidate variety. The maximum allowable frequency of each variant for each class of pedigreed seed must be given.

_____ **No deviant plants or variants have been observed during seed increase of this variety.**

CSGA standards should apply for off-types for each pedigreed seed class. _____

- 13.0** List the characteristics that are the most useful for distinguishing the candidate variety. Refer to the characteristics using the objective description key numbers.

- 14.0** Additional characteristics:

11-07-94

APPENDIX**METHODS AND ILLUSTRATIONS****3.1 Stem: fasciation**

The expression of fasciation varies considerably with environmental conditions, although the presence or absence of fasciation is usually clear.

3.2 Stem: vine length

The observations should be made on harvested plants at mature green seed stage. The measurement should include nodes with scale leaves. Both plant height at flowering and stem length at mature green seed stage may vary with site and season due to different responses to day length, temperature and soil moisture. Both characteristics are good discriminators within years at one site, however, and allow the separation of different varieties.

3.3 Stem: number of nodes up to and including the first flowering node

The expression can vary due to flower abortion under certain environmental conditions. Nodes with scale leaves should be included.

4.2 Leaf: average maximum number of leaflets

The maximum expression should be recorded over the whole plant. Although appearing to be continuously expressed, this characteristic is stable. Occasional plants may have a larger number of leaflets. The maximum number of leaflets for a sample of plants should be recorded and an average value calculated.


4.6 Leaf: dentation

The observations should be made over the whole plant, with the exception of the lowest six nodes and all aerial and basal branches.


PEA OBJECTIVE DESCRIPTION

APPENDIX


4.7 Leaf: degree of dentation



1 - very weak



3- weak



5 - medium



7 - strong



9 - very strong

PEA OBJECTIVE DESCRIPTION**APPENDIX****5.1 Stipule: development**

Rudimentary stipules are lanceolate and surface area is reduced significantly by up to 80%. Plants with 'Rabbit-eared' stipules are not examples of rudimentary stipules.

5.2 Stipule: size


The observations should be made at the second fertile node on stipules which have been detached from the plant and flattened.

5.5 Stipule: marbling

The observations should be made over the whole plant. Care has to be taken that foliage at the lowest nodes has not senesced before assessment. If assessed before flowering, the plant should have at least eight nodes, since flecking in some varieties may not be expressed at lower nodes.

5.6 Stipule: maximum density of marbling

The observations should be made over the whole plant.



PEA OBJECTIVE DESCRIPTION**APPENDIX****6.7 Flower: shape of base of standard**

The observations should be made on a sample from different plants. The standard should be detached and flattened on a hard surface and compared with example varieties before assigning a state.




3 - raised
(V-shaped)



5 - level
(straight)



7 -arched
(2 lobes)



9 - strongly arched

7.3 Pod: parchment

(1) The observation should be made on a sample from different plants when the pods are dry and papery.

(2) The pod should be opened along the suture without damaging the edges of the two valves. The distribution of sclerenchyma, which makes up the parchment, may either be observed by staining with Phoroglucinol in Hydrochloric Acid, or by reflecting light (preferably daylight) on the inside of the pod wall.

If parchment for any pod is difficult to determine, pods from other nodes on the same plant should be examined.

7.5 Pod: type of curvature

The observations should be made on the upper suture on a sample of plants. The maximum expression over the whole plant should be assessed. The 'hook end' on long podded types should be ignored when assessing curvature.

7.6 Pod: shape of distal part

The observations should be made only on varieties without thickened pod wall. They should be made on a sample of plants and on several nodes of each plant when pods are fully developed, but before any senescence. Care should be taken where pods are strongly curved, where the beak is longer than the pod tip, or where parchment is not entire. Some varieties have a blunt tip which is rounded, but the beak is higher up the pod.

PEA OBJECTIVE DESCRIPTION**APPENDIX****7.7 Pod: colour**

- (1) Each state should be treated separately.
- (2) Varieties with yellow pods may also have yellowish peduncles and sepals. In the presence of anthocyanin, colouration of the pods will appear red.
- (3) The appearance of green pods is the result of yellow, purple and blue-green colours not being expressed.
- (4) Blue-green pods are dark and slightly bluish, but not as blue as blue-green foliage. The colour develops with time and may be more accentuated in hotter, drier conditions.
- (5) The expression of purple pods can be variable and unstable, disappearing on the same plant or being reduced in its distribution on the pod.

8.0 DRY SEED CHARACTERISTICS

The provided seed should be mature and preferably not severely bleached, the assessment should be carried out within nine months after harvest. For varieties with anthocyanin pigment, tannins in the testa often darken with age, (usually after nine months) obscuring many characteristics. The observation is most clear under conditions of bright natural light; the assessment of some characteristics is difficult under artificial light.

8.1 Seed: shape of starch grain

- (1) After removing the testa, fine fragments of tissue should be extracted from the cotyledon and examined after having added water and been squashed gently between two microscope slides. Too much pressure during squashing results in fragmentation of the grains, too little pressure will not provide a layer thin enough for easy examination. This works best on pea flour (finely ground pea seed).
- (2) A microscope with transmitted light, using x16 eye-pieces and either x10 or x40 objectives, is most suitable for examination. For examination of compound grains, the larger objectives will be required.
- (3) Simple grains resemble wheat seeds or coffee beans in shape, often with what looks like a suture line running along their length.
- (4) Compound grains look irregularly star-shaped and appear to be made of a number of segments. The center of the grains may appear cross-shaped. Too much pressure during squashing causes fragmentation.

PEA OBJECTIVE DESCRIPTION**APPENDIX****8.2 Seed: colour of cotyledon**

The expression varies with environmental conditions:

- (i) bleaching, caused by sunlight or chemical changes in the plant, can remove colour from both green and yellow cotyledon seeds;
- (ii) colour becomes dull with age, even if seed is stored in cold, dark conditions;
- (iii) colour can darken in the presence of high amounts of Tragacanth oil occurring on the underside of the testa. This fades as the seed ages.

There is a range of colour from yellow to orange yellow and from pale to dark green.

8.3 Seed: black colour of hilum

- (1) The hilum colour can be influenced by the presence of tannin in the testa. If any loose tissue is present, the hilum area should be lightly polished with a cloth before recording.
- (2) Spontaneous mutation from melanin absent to melanin present can occur. This appears to be more prevalent in colored flowered types. The mutation rate is unknown.

8.4 Seed: shape

The shape can be influenced by environmental conditions, although it is generally consistent from year to year, provided the seed has reached its full development.

8.5 Seed: wrinkling of cotyledon

The observations should be made on harvested seed. 'Golf ball' and large dimples should be ignored as these can also be found on smooth seeded (non-wrinkled) types. Cylindrically shaped seed types should be assessed carefully, because some are smooth seeded.

8.6 Seed: size

The observations should be made on harvested seed only. The weight varies markedly from site to site but can be useful as a discriminator; it varies to a lesser extent from season to season at one site. Immature and infected seeds should be excluded; the seed should be dry (approximately 10-15% moisture content) at time of recording.

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE**EXHIBIT E**
STATEMENT OF THE BASIS OF OWNERSHIP

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

1. NAME OF APPLICANT(S) SVALÖF WEIBULL AB	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER SW 975504	3. VARIETY NAME SW CIRCUS
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country) S-268 81 SVALÖV, SWEDEN	5. TELEPHONE (include area code) 46-418-667000	6. FAX (include area code) 46-418-667100
7. PVPO NUMBER 200200115		
8. Does the applicant own all rights to the variety? Mark an "X" in appropriate block. If no, please explain. <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO		

9. Is the applicant (individual or company) a U.S. national or U.S. based company? If no, give name of country SWEDEN		<input type="checkbox"/> YES <input checked="" type="checkbox"/> NO
10. Is the applicant the original owner? <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO If no, please answer <u>one</u> of the following:		
a. If original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. national(s)? <input type="checkbox"/> YES <input type="checkbox"/> NO If no, give name of country		
b. If original rights to variety were owned by a company(ies), is(are) the original owner(s) a U.S. based company? <input type="checkbox"/> YES <input type="checkbox"/> NO If no, give name of country		
11. Additional explanation on ownership (if needed, use reverse for extra space):		

PLEASE NOTE:

Plant variety protection can be afforded only to owners (not licensees) who meet one of the following criteria:

1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
2. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definition.

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

STD-470-E (07-97) (Destroy previous editions).


Electronic version designed using WordPerfect InForms by USDA-AMS-IMB.

200200115

U.S. Plant Variety Protection Act - Authorization of an Agent

I hereby authorize Bonis & Company Ltd. to act, for all purposes of this Act, on behalf of me as my agent for the Field Pea Variety, **SW CIRCUS**.

Signature:

.....

Howard K. Love
Canadian Research Director
Svalof Weibull AB

Date:

2002-03-11

Address:

2-411 Downey Road
Saskatoon, Sask.
S7N 4L8
Canada